

Acta Crystallographica Section C

Crystal Structure Communications

Volume 50, Part 5 (May 1994)

organic compounds

[pdf](#) [cif](#) [3d view](#) [similar papers](#) [contents of issue](#) [buy](#)

Acta Cryst. (1994). **C50**, 771-773 [doi:10.1107/S0108270193010741]

5,6-Dihydroxy-7-methoxyflavone

M. Shoja

[download](#) bibliographic record in [BIBTeX](#) [format](#)

This is the first page of the item you requested.

C(17)-C(18)-C(19)	0.9951 (2)	0.5679 (4)	-0.1656 (2)	0.9443 (13)
C(18)-C(19)-C(20)	0.9079 (2)	0.5459 (3)	-0.0336 (3)	0.0514 (18)
C(19)-C(20)-C(21)	0.8023 (3)	0.5431 (3)	0.0920 (3)	0.0490 (17)
C(20)-C(21)-C(22)	0.1361 (2)	0.2539 (4)	0.1591 (2)	0.0412 (13)
C(21)-C(22)-C(23)	0.1985 (3)	0.4022 (4)	0.3742 (2)	0.0445 (16)
C(22)-C(23)-C(24)	0.1322 (2)	0.5159 (5)	0.4718 (3)	0.0450 (18)
C(23)-C(24)-C(25)	0.1489 (3)	0.4832 (5)	0.5337 (2)	0.0546 (19)
C(24)-C(25)-C(26)	0.2276 (4)	0.7064 (3)	0.4611 (3)	0.0681 (23)
C(25)-C(26)-C(27)	0.2040 (2)	0.7311 (5)	0.4644 (3)	0.0593 (22)
C(26)-C(27)-C(28)	0.2799 (4)	0.5664 (3)	0.3321 (3)	0.0606 (23)

Table 2. Selected geometric parameters (\AA , $^\circ$)

P-C(1)	1.815 (2)	P-C(20)	1.819 (2)
P-C(23)	1.835 (3)	C(1)-C(10)	1.391 (4)
C(10)-C(11)	1.479 (4)	C(11)-C(20)	1.391 (4)
H(81)-H(81)	2.33 (5)		
P-C(1)-C(2)	125.7 (2)	P-C(1)-C(10)	153.5 (1)
P-C(20)-C(1)	114.0 (2)	P-C(20)-C(10)	105.9 (2)
P-C(21)-C(22)	119.8 (2)	P-C(21)-C(26)	131.7 (3)
C(1)-P-C(20)	89.4 (3)	C(1)-P-C(21)	109.9 (1)
C(10)-P-C(21)	108.2 (1)	C(2)-C(1)-C(10)	121.8 (2)
C(1)-C(10)-C(9)	115.0 (2)	C(1)-C(10)-C(20)	112.4 (2)
C(9)-C(10)-C(11)	120.6 (3)	C(10)-C(11)-C(2)	129.4 (3)
C(10)-C(11)-C(20)	112.6 (2)	C(1)-C(11)-C(20)	118.1 (2)

Table 3. Dihedral angles ($^\circ$) between least-squares planes

Plane 1: P, C(3), C(10), C(11), C(20). Plane 2: C(1)-C(4), C(9), C(10).			
Plane 3: C(4)-C(9). Plane 4: C(11), C(12), C(17)-C(20). Plane 5: C(32)-C(17). Plane 6: C(21)-C(26). Nap1: C(1)-C(10), Nap2: C(11)-C(20).			
Plane 1-Plane 2	0.07 (2)	Plane 1-Plane 4	12.03 (3)
Plane 1-Plane 6	92.9 (2)	Plane 2-Plane 3	12.2 (2)
Plane 2-Plane 4	18.7 (2)	Plane 3-Plane 5	38.2 (2)
Plane 4-Plane 5	9.8 (1)	Nap1-Nap2	28.7 (3)

The positional parameters for all non-H atoms were determined by direct methods (Sheldrick, 1983). The refinements were carried out by full-matrix least-squares techniques (Imoto, 1990). All H atoms were located in a difference Fourier map. The $\Delta\rho^+$ and $\Delta\rho^-$ components of anomalous dispersion were included in the calculation for the P atom (Cromer & Ibers, 1974). The refined structure showed R and wR values of 0.0494 and 0.0601. Thus, we decided that the title compound has the absolute configuration P . All calculations were carried out on a NEC ACOS 930S computer at the Research Center for Protein Engineering, Institute for Protein Research, Osaka University.

The authors thank Dr Hideo Imoto for the least-squares program (ANYBLK). Financial support from CIBA GEIGY Foundation (Japan) for the Promotion of Science is gratefully acknowledged.

Lists of structure factors, anisotropic displacement parameters, H -atom coordinates and complete geometry have been deposited with the British Library Document Supply Centre as Supplementary Publication No. SUP 11696 (37 pp.). Copies may be obtained through The Technical Editor, International Union of Crystallography, 3 Abbey Square, Chester CH1 3HL, England. [CIF reference: A51077]

References

- Cromer, D. Y. & Ibers, J. A. (1974). *International Tables for X-ray Crystallography*, Vol. IV, Table 2.3.1. Birmingham, Kynoch Press. (Present distributor Kluwer Academic Publishers, Dordrecht.)

© 1994 International Union of Crystallography
Printed in Great Britain. All rights reserved.

Imoto, H. (1990). *ANYBLK*. Program for Least-Squares Refinement. Department of Chemistry, Univ. of Tokyo, 015, Japan.

Johnson, C. K. (1976). *ORTEP*. Report ORNL-5138, Oak Ridge National Laboratory, Tennessee, USA.

Kagan, H. B. (1985). *Asymmetric Catalysis*, Vol. 5, edited by J. D. Morrison, pp. 1-6. New York: Academic Press.

Nosch, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst.* **A24**, 351-359.

Parsons, W. A., RajanBabu, T. V. & Burk, M. J. (1993). *Science*, **259**, 479-483.

Sheldrick, G. M. (1985). *SHELX586*. Program for the Solution of Crystal Structures. Univ. of Göttingen, Germany.

Tom, K., Tashiro, H. & Yamagata, T. (1992). *Abstracts of Symposium on Organometallic Chemistry, Japan*, pp. 158-159.

Watson, A. A., Wells, A. C. & Wild, S. B. (1993). *J. Organomet. Chem.* **445**, 71-78.

Acta Cryst. (1994), **C50**, 771-773

5,6-Dihydroxy-7-methoxyflavone

M. SHOJA

Chemistry Department, Fordham University, Bronx, NY 10458, USA

(Received 5 March 1993; accepted 5 October 1993)

Abstract

The 5-hydroxy group of the title compound, 5,6-dihydroxy-7-methoxy-2-phenyl-4*H*-1-benzopyran-4-one, $C_{15}H_{10}O_5$, forms a cyclic intramolecular hydrogen bond $O(3)\cdots H-O(2)$, $H\cdots O \approx 1.71$ (3) Å, with the carbonyl group. The heterocyclic ring is not coplanar with the phenyl ring. The C(7) methoxy group is in the plane of the γ -benzopyrone ring with the torsion angle $C(11)\cdots O(4)\cdots C(7)\cdots C(8) \approx 2.9$ (3) $^\circ$.

Comment

The dihedral angle of 12.2 (2) $^\circ$ between the phenyl ring and the γ -benzopyrone portion of the molecule (I) is significantly different from those of two related structures, 5-hydroxy-7-methoxyflavone (Shoja, 1989) and 5-hydroxyflavone (Shoja, 1990), with dihedral angles of 24.8 (2) and 5.7 (7) $^\circ$, respectively. Given the wide range of dihedral angles and the fact that all three of these molecules contain hydrogen bonding, it is unlikely that any planarity is solely a



Acta Crystallographica Section C
ISSN 0108-2701 © 1994

Find reference: Volume Page

Search: From to [Advanced search](#)

Copyright © International Union of Crystallography
IUCr Webmaster